

Research paper

Perceived emotional social support in bereaved spouses mediates the relationship between anxiety and depression



Nicholas C. Jacobson*, Kayla A. Lord, Michelle G. Newman

The Pennsylvania State University, United States

A B S T R A C T

Background: Prior research has shown that anxiety symptoms predict later depression symptoms following bereavement. Nevertheless, no research has investigated mechanisms of the temporal relationship between anxiety and later depressive symptoms or examined the impact of depressive symptoms on later anxiety symptoms following bereavement.

Methods: The current study examined perceived emotional social support as a possible mediator between anxiety and depressive symptoms in a bereaved sample of older adults ($N = 250$). Anxiety and depressive symptoms were measured at Wave 1 (immediately after bereavement), social support was measured at Wave 2 (18 months after bereavement), and anxiety and depressive symptoms were also measured at Wave 3 (48 months after bereavement).

Results: Using Bayesian structural equation models, when controlling for baseline depression, anxiety symptoms significantly positively predicted depressive symptoms 48 months later. Further, perceived emotional social support significantly mediated the relationship between anxiety symptoms and later depressive symptoms, such that anxiety symptoms significantly negatively predicted later emotional social support, and emotional social support significantly negatively predicted later depressive symptoms. Also, when controlling for baseline anxiety, depressive symptoms positively predicted anxiety symptoms 48 months later. However, low emotional social support failed to mediate this relationship.

Conclusions: Low perceived emotional social support may be a mechanism by which anxiety symptoms predict depressive symptoms 48 months later for bereaved individuals.

1. Introduction

Bereavement and spousal loss is especially common in later life with 24.7% of adults ages 65 and older being widowed (United States Census Bureau, 2015). One study found that over a course of 2.5 years, 8.1% of older adults reported losing a spouse (Williams et al., 2007). Bereavement also was associated with both anxiety (MacCallum et al., 2015; Meier et al., 2013; Prigerson et al., 1996; Byrne and Raphael, 1997; Kreicbergs et al., 2004; Valdimarsdottir et al., 2004; Mitchell et al., 2009) and depressive symptoms (Kreicbergs et al., 2004; Mitchell et al., 2009; Stroebe et al., 2005, 1996; Murphy, 1988; Jozwiak et al., 2013; Ghesquiere et al., 2013). Additionally in those experiencing bereavement, higher anxiety was associated with higher perceived loss of control (Ong et al., 2005), lower energy (Boelen and Prigerson, 2007), increased suicidal ideation (Chen et al., 1999), increased risk of a heart attack (Chen et al., 1999), increased risk of stomach problems (Chen et al., 1999), poorer health (Boelen and Prigerson, 2007), and greater difficulty sleeping (Boelen and Prigerson,

2007) compared to those with lower anxiety. Similarly, in those experiencing bereavement, higher depression was associated with higher blood pressure (Chen et al., 1999), poorer health (Chen et al., 1999), greater cognitive impairment (Ward et al., 2007) and poorer coping (Bennett et al., 2005) compared to those with lower depression. Given the wide-ranging influence of anxiety and depressive symptoms in those who have experienced loss, it is important to understand the longitudinal relationship between anxiety and depressive symptoms following bereavement as well as potential mediators of this relationship.

In evidence on the longitudinal relationship between anxiety and depression outside the context of bereavement, a meta-analysis examining 29 studies and over 20,000 people, showed that anxiety symptoms positively predicted later depressive symptoms with an estimated longitudinal association of $r = 0.34$ (Jacobson and Newman, 2012b; van Baarsen, 2002). Likewise, anxiety symptoms predicted depressive symptoms across many time scales, ranging from hours to years (Jacobson and Newman, 2012a, 2014, 2016; Starr and Davila,

* Correspondence to: Pennsylvania State University, 378 Moore Building, University Park, PA 16802, United States.
E-mail address: njacobson@psu.edu (N.C. Jacobson).

2012a, 2012b; Swendsen, 1997; Starr et al., 2016). Similarly, in a meta-analysis of the broader literature including approximately 8000 persons, depressive symptoms predicted later anxiety symptoms with an average longitudinal association of $r = 0.35$ (Jacobson and Newman, 2012b; Jacobson and Newman, in preparation).

Nonetheless, literature on the longitudinal relationship between anxiety and depression in the context of bereavement is much more scarce. Following bereavement, two studies found that anxiety symptoms predicted later depressive symptoms (Prigerson et al., 1996; Boelen and Prigerson, 2007). At the same time, the relationship between depression and later anxiety has not been examined within this context. Thus, despite robust bi-directional associations between anxiety and depression in younger populations, no studies have actually tested whether a bi-directional relationship exists between anxiety and depression in older adult populations. In addition, no prior research has examined the mechanism of this relationship in the context of bereavement or in an elderly population. Studying such mechanisms within older adults will become increasingly important as, by the year 2033, those who are 65-years and older will outnumber those 18 and younger in the United States (Colby and Ortman, 2015).

There is reason to focus on perceived emotional social support as a possible mediator of this relationship. Following widowhood, the feeling of anxiety and dread about other bad things happening in the future is theorized to lead many either to emotionally and socially withdraw to conserve one's energy for personal coping (Hogan et al., 1996), or to engage in behaviors that might push others away. The impact of such withdrawal or negative behaviors may be greater in aging widows/ widowers, as older adulthood (even outside the context of bereavement) is a time frequently associated with decreasing social support, due to increasing physical barriers and loss in one's social circles (Mullins and Dugan, 1990). Thus, bereavement in older adulthood is likely to be a time of considerable anxiety, dread, and tension, and these feelings are likely to lead to greater withdrawal, poor emotional disclosure, and unaffiliative behaviors.

A lack of emotional support following expressions of anxiety may be particularly impactful among older adults. Low social support in this population may contribute to feelings of loneliness, isolation (van Baarsen, 2002; Golden et al., 2009; Van Baarsen et al., 1999), and emotional isolation (van Baarsen, 2002). By not having sufficient support and outlets to cope with and process the loss, social and emotional isolation may be a primary mechanism between bereavement anxiety and later depression (Cornwell and Waite, 2009). In sum, high anxiety following widowhood is theorized to lead to friends' departure, low emotional support, isolation, and ultimately depression.

Some evidence for this theory comes from nonbereaved younger samples. Higher anxiety symptoms were associated with more difficulty in the social domain [including difficulty making friendships; (Langston and Cantor, 1989; Scharfstein et al., 2011)] and more difficulty with emotional self-disclosure (Kahn and Garrison, 2009; Post et al., 1978). Chronic worrying also has been associated with impacting significant others in unaffiliative ways (Erickson et al., 2016). In longitudinal studies, anxiety symptoms unidirectionally predicted later low social support, and such lack of social support did not increase subsequent anxiety symptoms (Friedmann et al., 2014; Paterson et al., 2015). In addition, lower levels of perceived emotional social support predicted higher depressive symptoms longitudinally (Stroebe et al., 2005, 1996; Aneshensel and Frerichs, 1982; Cohen and Hoberman, 1983) following negative life events. There is also evidence that perceived emotional social support may be a mechanism by which anxiety leads to depression. One prior study found that perceptions of intimacy in close and group relationships mediated the longitudinal relationship between anxiety and later depressive symptoms (Jacobson and Newman, 2016). In another study, interpersonal dysfunction mediated the relationship between generalized anxiety disorder and depressive symptoms (Starr et al., 2014).

Additional evidence comes from cross-sectional data using bereavement samples. For example, anxiety symptoms were associated with

low emotional social support in persons who had experienced a stillbirth [which has been noted as a form of bereavement; (Cacciatore et al., 2009; Condon, 1986)] and in bereaving spouses (Bergman et al., 2010). In addition, in observations of widows discussing their emotional reactions to the loss of their spouses, those perceived as less well-adjusted evoked greater frustration and less compassion from onlookers, compared to those who were perceived as better adjusted (Keltner and Bonanno, 1997). In turn, deficits in support networks, loneliness, and the amount of emotional support received from one's friendships have also been cross-sectionally linked to depression in older adults experiencing bereavement (Prince et al., 1997; van Grootheest et al., 1999). Further, among the elderly experiencing bereavement, a strong relationship has been found between lacking help in making decisions and feeling more blue or depressed (Balaswamy et al., 2004). Thus, persons experiencing and expressing their anxiety may elicit low compassion and a lack of emotional social support from others, leading to fewer social resources, lower emotional support, less helping making decisions and ultimately depression.

The goal of the current study was to examine perceived emotional social support as a mechanism of the relationship between post-bereavement earlier anxiety and later depressive symptoms utilizing a longitudinal design in an older adult sample. We also tested whether the relationship between anxiety and depression was unidirectional or bidirectional following bereavement in older adults. Face-to-face interviews were conducted 6-months (Wave 1), 18-months (Wave 2), and 48-months (Wave 3) after a spouse's death. A body of literature regarding bereavement trajectories supports this timing of measurement points. The precedent in spousal bereavement literature is to assess symptoms post bereavement at 3–7 months, then between 12 and 18 months, and at multiple time points in years with a particular interest in the first five years after bereavement (Utz et al., 2011, 2014; Bonanno et al., 1999; Middleton et al., 1996; Field et al., 2003; Sasson and Umberson, 2014; Galatzer-Levy and Bonanno, 2012; Holland et al., 2014). Some research shows that bereavement symptoms are still present decades after the loss (Carnelley et al., 2006). The interviews consisted of a large set of questions including validated anxiety and depression scales. Social support was defined using two highly correlated items. Based on previous research, we hypothesized that: (1) anxiety symptoms (at Wave 1) would positively predict depressive symptoms (Wave 3) three and half years later while controlling for baseline depressive symptoms (Wave 1), (2) depressive symptoms (Wave 1) would positively predict anxiety symptoms (Wave 3) three and a half years later while controlling for baseline anxiety symptoms (Wave 1), (3) emotional social support (Wave 2) would mediate the relationship between anxiety symptoms (Wave 1) and later depressive symptoms (Wave 3), such that anxiety symptoms would negatively predict emotional social support, and emotional social support would negatively predict depressive symptoms, and (4) emotional social support (Wave 2) would not significantly mediate the relationship between depression (Wave 1) and later anxiety (Wave 3).

2. Method

2.1. Participants

Participants ($N = 250$) were collected through the Changing Lives of Older Couples (CLOC): A Study of Spousal Bereavement in the Detroit Area public use data set (Nesse et al., 2006). Data collection took place in three waves between 1987 and 1993. Individuals were identified through a two-stage area probability sample (Nesse et al., 2006). To be eligible for the study, participants needed to speak English, be at least 65 years old, and married at the time of initial contact. Following initial contact, death records of the state of Michigan were monitored, and participants were contacted 6-, 18-, and 48-months following the loss of their spouse (Nesse et al., 2006). Data were collected through

Table 1
Anxiety, Depression, and Social Support Scales.

Measure	Item	Scale	
SCL-90-R	[How much have you been bothered by] nervousness or shakiness?	Not at all 1–5 extremely	
	...trembling?	Not at all 1–5 extremely	
	...feeling suddenly scared for no reason?	Not at all 1–5 extremely	
	...feeling fearful?	Not at all 1–5 extremely	
	...heart pounding or racing?	Not at all 1–5 extremely	
	...feeling tense and keyed up in the past seven days?	Not at all 1–5 extremely	
	...spells of terror or panic?	Not at all 1–5 extremely	
	...feeling so restless you couldn't sit still?	Not at all 1–5 extremely	
	...feeling that something bad is going to happen to you?	Not at all 1–5 extremely	
	...thoughts and images of a frightening nature?	Not at all 1–5 extremely	
	CES-D	I felt depressed.	Hardly ever 1–3 most of the time
		I felt that everything I did was an effort.	Hardly ever 1–3 most of the time
		My sleep was restless.	Hardly ever 1–3 most of the time
I was not happy.		Hardly ever 1–3 most of the time	
I felt lonely.		Hardly ever 1–3 most of the time	
People were unfriendly.		Hardly ever 1–3 most of the time	
I did not enjoy life.		Hardly ever 1–3 most of the time	
I did not feel like eating. My appetite was poor.		Hardly ever 1–3 most of the time	
I felt sad		Hardly ever 1–3 most of the time	
I felt that people disliked me.		Hardly ever 1–3 most of the time	
I could not get 'going.'		Hardly ever 1–3 most of the time	
Emotional Social Support	On the whole, how much do your friends make you feel loved and cared for?	A great deal 1–5 not at all	
	How much are your friends and relatives willing to listen when you need to talk about your worries or problems?	A great deal 1–5 not at all	

Note: This table represents the items from the Symptoms Checklist 90-Revised, the Center for Epidemiological Studies Depression Scale, and the constructed Social Support Scale.

interviews. Wave 1 interviews were conducted 6-months after the loss of a spouse ($N = 250$; 86% female, M age =70.11, 84.4% Caucasian, 15.6% African American). Wave 2 interviews were conducted 18-months post-bereavement ($N = 210$), and the third and final wave of data was collected 48-months post-bereavement ($N = 106$).

2.2. Measures

2.2.1. Anxiety subscale of the Symptom Checklist-90 Revised

Anxiety symptoms were measured during each wave with the 10-item anxiety subscale of the Symptom Checklist-90 Revised (SCL-90-R) (Derogatis, 1983), which is a self-report questionnaire designed to measure anxiety symptoms in the general population. See Table 1 for items. The 10-item anxiety subscale has high convergent validity with the Minnesota Multiphasic Personality Inventory (MMPI) anxiety subscale (Derogatis et al., 1976) and the Beck Anxiety Inventory (BAI) (Steer et al., 1993). Furthermore, high levels of invariance across gender have been noted (Derogatis and Cleary, 1977a), and high construct validity has been found as well (Derogatis and Cleary, 1977b). In the current sample, the internal consistency of this subscale was high ($\alpha = 0.84$) (Nesse et al., 2006). Responses were measured on a five point Likert scale from 1 (not at all) to 5 (extremely). (Table 2).

Table 2
Scale Correlations and Descriptive Statistics.

	1	2	3	4	5
1. Anxiety Wave 1	1				
2. Depression Wave 1	0.40	1			
3. Emotional Social Support Wave 2	0.29	0.08	1		
4. Anxiety Wave 3	0.23	0.20	0.10	1	
5. Depression Wave 3	0.35	0.33	0.23	0.38	1
<i>M</i>	12.99	11.04	3.74	11.16	10.44
<i>SD</i>	4.11	1.62	1.83	2.41	1.15
Range	10–31	6–17	2–10	10–27	6–15

Note: This table first presents correlations between items in the first five rows of the table. Following this, the mean, standard deviation, and range of each of the scale measures.

2.2.2. Center for Epidemiological Studies Depression Scale

The CES-D, a self-report questionnaire meant to evaluate depressive symptoms in the general population (Radloff, 1977), was delivered during all 3 waves. The original scale had 20-items, however, an 11-item version was used here. See Table 1 for items. The 11-item version was deemed preferable for this study because the full scale was found to be taxing for older adults (Kohout, 1992). Responses were measured on a three point Likert scale from 1 (hardly ever) to 3 (most of the time). The 20-item scale showed good convergent validity when compared to the SCL-90-R depression scale ($r = 0.73–0.89$) (Weissman et al., 1977), the BDI, and the MMPI-II (Bush et al., 2004). The 11-item subscale also has been found to have good construct validity (Gellis, 2010). In the current sample, internal consistency of this scale was high ($\alpha = 0.84$) (Nesse et al., 2006).

Nevertheless, the 11-item CES-D contains questions that are not specific to depressive symptoms. For example, it also measures social relationships (i.e. people being unfriendly, people disliking them, and loneliness) and symptoms that overlap with anxiety (i.e. restlessness, appetite disturbance). To ensure that these items did not undermine the theoretical constructs, they were removed for the primary analyses (note that additional analyses were also conducted with the original 11-item CES-D; see planned analyses below). The remaining 6-item version still had high internal consistency ($\alpha = 0.80$).

2.2.3. Perceived emotional social support scale

Two items were used to measure perceived emotional social support: (1) “On the whole, how much do your friends make you feel loved and cared for?” and (2) “How much are your friends and relatives willing to listen when you need to talk about your worries or problems?” This scale was given during Wave 2, 18-months after the loss of a spouse. Internal consistency of this scale was high ($\alpha = 0.80$) (Nesse et al., 2006). Responses were measured on a five point Likert scale from 1 (a great deal) to 5 (not at all). Items were then reverse coded, such that higher scores indicated higher levels of social support.

Although this scale was developed for this study specifically, it shares a great deal of overlap between previously validated scales. In particular, it is similar to (1) The Medical Outcomes Study (MOS) Social Support [which likewise assesses social support using items such as “love and affection”, “listen to you”, “confide in”, and “share worries

with”; (Sherbourne and Stewart, 1991)]; (2) The Duke-UNC Functional Social Support Questionnaire [also using the following similar items “Chances to talk with someone about [all types] of problems” and “Love and affection” (Broadhead et al., 1988)], and (3) the Interview Schedule for Social Interaction [similarly to the current study it asks “Do you have someone you can share your most private feelings with (confide in) or not?” as well as multiple items assessing the perceived love and care (Henderson et al., 1980; Uden and Orth-Gomer, 1989)]. Thus, although we used a novel two-item scale, it contained high item content validity with previously validated scales.

2.2.4. Covariates

Note that the analyses were repeated with and without controlling for the following sources of stress. During the second wave, persons were asked if they (1) experienced a life-threatening illness or injury, (2) have a serious but not life-threatening illness, (3) were robbed or burglarized, (4) involuntarily lost a job for reasons other than retirement, (5) have serious financial problems or difficulties, and (6) moved to a new residence.

2.3. Planned analyses

All primary analyses used Bayesian structural equation models (BSEM) in Mplus 7. BSEM has unique advantages over frequentist SEM: (1) unlike frequentist SEM, which usually requires considerably larger sample sizes, BSEM coverage rates have been shown to perform well in simulation studies with sample sizes ranging from 25 to 1000 (Yuan and MacKinnon, 2009; Gelman and Rubin, 1992), (2) BSEM does not have problems with non-convergence, nonsensical values, and Heywood cases (Enders, 2010; Rupp et al., 2004), and (3) BSEM tends to perform well under conditions of multicollinearity (Winship, 1999; Jagpal, 1982). Notably, BSEM also handles ordinal data well, allowing distributions to be directly set within the modeling framework (Asparouhov and Muthen, 2010). Missing data was handled through random forest multiple imputation using the missForest package, which has greater accuracy than other multiple imputation methods (Stekhoven and Bühlmann, 2012).

Prior to the primary analyses, to ensure that there were not baseline differences between completers and non-completers, a Bayesian independent samples *t*-test was conducted on the factor scores of the anxiety and depression variables at Wave 1 using JASP (Team, 2016). Note that the test statistic in a Bayesian *t*-test is the Bayes factor. A Bayes factor is a model-based estimate of the odds of the alternative hypothesis being true compared to the null hypothesis. A Bayes Factor ranges from 0 to infinity, and the higher the Bayes factor the more evidence there is to reject the null hypothesis (i.e. that there are no differences between completers and non-completers). Values less than 1 suggest that there is more support for the null-hypothesis than there is for the alternative hypothesis (Kass and Raftery, 1995). Note that the default non-informative Cauchy prior was used (Team, 2016).

First, we tested the first and second hypotheses that anxiety symptoms at Wave 1 would predict depressive symptoms at Wave 3, controlling for Wave 1 anxiety symptoms, and depressive symptoms at Wave 1 would predict anxiety symptoms at Wave 3, controlling for Wave 1 depression symptoms. Item residuals between Wave 1 and Wave 3 were allowed to covary. Next, we tested the hypothesis that the relationship between anxiety symptoms at Wave 1 and depressive symptoms at Wave 3 would be mediated by perceived emotional social support, controlling for Wave 1 depression symptoms (we also tested social support mediating the relationship between depression and later anxiety, controlling for earlier anxiety). Mediation was estimated directly using indirect and total effect model parameters, and these parameters were estimated directly using the Markov Chain Monte Carlo (MCMC) estimates. As with other mediation approaches (i.e. bootstrapping), BSEM parameter estimates do not make any distributional assumptions of the parameters in the estimation (Yuan and MacKinnon, 2009).

Model fit was assessed based on the posterior predictive p-value (PPP) (Muthén and Asparouhov, 2012a, 2012b), which uses the likelihood-ratio chi-square test of the estimated data against the observed data. Based on simulation work, PPP values above 0.050 represent good model fit (Asparouhov and Muthen, 2010; Muthén and Asparouhov, 2012a). Instead of p-values, which have been shown to be highly unreliable (Cumming, 2014), BSEM uses Bayesian credible intervals wherein the model suggests that the true parameter has a 95% chance of falling within the specified credible interval bounds (Newcombe, 1998) and is determined to be significant if credible intervals do not contain 0. Non-informative priors were used in the modeling approach as recommended by Asparouhov and Muthen (Asparouhov and Muthen, 2010) and Muthén and Asparouhov (Muthén and Asparouhov, 2012a). All regression coefficients are also presented with a Cohen's *d* effect size based on the following formula, $d = \frac{\beta}{SD(\beta)} * \sqrt{\frac{2}{n}}$ (Dunlap et al., 1996), and consequently the magnitude of the effect size suggests that 0.2 represents a small effect, 0.5 represents a medium effect, and over 0.8 represents a large effect (Cohen, 1988).

All primary analyses were conducted using the CES-D items that did not overlap with emotional social relationships or with anxiety (see CES-D section above). However, to be comprehensive, all analyses were repeated with the 11-item CES-D scale. Further, to ensure that the results were not driven by experiencing other sources of stress, all analyses were repeated using each stress variable as a covariate in predicting anxiety and depression at wave 3. Note that both sets of results are reported in footnotes below.

3. Results

3.1. Differences between completers and non-completers

For baseline anxiety, the results suggested that it was unlikely that there was a difference (BF = 0.764) between completers ($M = 12.370$, $SD = 3.070$) and non-completers ($M = 13.380$, $SD = 4.647$). Likewise, for depression the results suggested that it was unlikely that there was a difference (BF = 0.144) between completers ($M = 11.04$, $SD = 1.707$) and non-completers ($M = 11.00$, $SD = 1.707$).

Hypotheses 1–2 : Anxiety Predicts Later Depression and Depression Predicts Later Anxiety.

The first analysis examined our first and second hypotheses. The model demonstrated good fit (PPP = 0.100). Supporting our first hypothesis, anxiety symptoms significantly positively predicted subsequent depressive symptoms when controlling for baseline depression symptoms ($B = 0.331$, $CI = 0.004–0.682$, $d = 0.271$; see Fig. 1).^{1,2} Likewise, supporting our second hypothesis, depressive symptoms significantly predicted later anxiety symptoms when controlling for baseline depression symptoms ($B = 0.330$, $CI = 0.027–0.632$, $d = 0.301$).

Hypotheses 3–4 : Emotional Social Support as a Mediator.

The next set of analyses examined our third and fourth hypotheses. The model demonstrated good fit (PPP = 0.058). The total effect of anxiety at Wave 1 on depression at Wave 3 was significant ($B = 0.368$, $CI = 0.009–0.774$, $d = 0.274$), suggesting that anxiety continued to predict later depression when emotional social support was included in

¹ Note that the model including the 11-item CES-D also had good fit (PPP = 0.058). Also, the pattern of anxiety predicting depression was consistent with the primary findings, as anxiety also significantly ($B = 0.387$, $CI = 0.079–0.737$, $d = 0.330$) positively predicted later depression in using the 11-item CES-D. Likewise, the results also showed that the model using the 11-item CES-D was consistent in that depression at wave 1 significantly predicted anxiety at wave 3 ($B = 0.468$, $CI = 0.154–0.873$, $d = 0.361$).

² Note that the model including the stress covariates also had good fit (PPP = 0.058). Also, the pattern of anxiety predicting depression was consistent with the primary findings, as anxiety also significantly ($B = 0.410$, $CI = 0.807–0.387$, $d = 0.290$) positively predicted later depression while controlling for later stress. Likewise, the results also showed that the model using the stress covariates was consistent in that depression at wave 1 significantly predicted anxiety at wave 3 ($B = 0.444$, $CI = 0.115–0.780$, $d = 0.361$).




Fig. 1. This figure depicts the first model results where anxiety and depressive symptoms at wave 1 predict anxiety and depressive symptoms at wave 3. Solid lines represent significant connections, whereas dotted lines represent insignificant connections.




Fig. 2. This figure depicts the second model results where social support at wave 2 mediates the relationship between anxiety at wave 1 and depressive symptoms at wave 3. Note that this figure also contains the relationship between anxiety and depression at wave 1 predicting anxiety and depression at wave 3. Solid lines represent significant connections, whereas dotted lines represent insignificant connections.

the model. Supporting our third hypothesis, anxiety symptoms significantly ($B = -0.288, CI = -0.587 - -0.005, d = -0.275$) negatively predicted later emotional social support, and social support significantly ($B = -0.496, CI = -0.711 - -0.297, d = -0.662$) negatively predicted later depressive symptoms (see Fig. 2). Further, the indirect effect of anxiety on depression through emotional social support was significantly positive ($B = 0.138, CI = 0.003 - 0.308, d = 0.253$), suggesting that perceived emotional social support significantly mediated the relationship between anxiety symptoms and later depressive symptoms. Additionally, the direct relationship between anxiety symptoms and later depressive symptoms was no longer significant ($B = 0.227, CI = -0.122 - 0.609, d = 0.176$). Based on the indirect and total effect estimates, emotional social support mediated 38% of the total variance

in anxiety predicting later depression. These results suggest that low emotional social support mediates a substantial portion of the variation between early anxiety and later depression.

Supporting our fourth hypothesis, in contrast to social support significantly mediating the relationship between anxiety and later depression, the relationship between earlier depression and later social support was not significant ($B = -0.118, CI = -0.405 - 0.158, d = -0.143$), and likewise, social support did not significantly predict later anxiety ($B = -0.055, CI = -0.247 - 0.137, d = -0.080$). Although the total effect of depression on later anxiety remained significant with social support in the model ($B = 0.356, CI = 0.057 - 0.698, d = 0.311$), the indirect effect of depression on later anxiety mediated by social support was not significant ($B = 0.003, CI = -0.003 - 0.051, d = 0.022$). Thus,

social support uniquely mediated the relationship between anxiety and later depression, and not vice versa.^{3,4}

4. Discussion

Supporting prior research using bereaved samples (Prigerson et al., 1996; Boelen and Prigerson, 2007) and the wider literature of anxiety and later depressive symptoms (Jacobson and Newman, 2012a, 2012b, 2014, 2016; Starr and Davila, 2012a, 2012b; Swendsen, 1997), the current results indicated that higher anxiety symptoms predicted higher depressive symptoms 48-months post bereavement. Additionally, depression following bereavement predicted anxiety symptoms 48-months later, suggesting a bi-directional relationship between anxiety and depressive symptoms post-bereavement. Furthermore, higher anxiety symptoms predicted lower emotional social support and lower emotional support predicted higher depressive symptoms. Thirty-eight percent of the relationship between earlier anxiety symptoms and later depressive symptoms was explained by the latter analysis. Nonetheless, perceived emotional social support failed to mediate the relationship between depression and later anxiety.

Findings in this age group may be particularly noteworthy given the unique developmental contexts of older adulthood. Older adults are more likely to experience declining physical health (Sibbritt et al., 2007), bereavement (Hansson and Stroebe, 2007), greater social isolation (Dykstra et al., 2005), and a loss of social support (Jensen et al., 2014). During such a pivotal time, anxious reactions to spousal loss are common (Maccallum et al., 2015; Meier et al., 2013; Prigerson et al., 1996; Byrne and Raphael, 1997; Kreicbergs et al., 2004; Valdimarsdottir et al., 2004; Mitchell et al., 2009), as the loss of one's spouse often signifies the beginning of feeling socially isolated (Mullins and Dugan, 1990; Victor et al., 2000). Supporting prior theories that anxiety reactions to bereavement in older adulthood lead to social withdrawal to conserve one's energy for personal coping (Hogan et al., 1996), and contextualizing findings that less well-adjusted reactions to grief evoke frustration and low compassion from others (Keltner and Bonanno, 1997), the current findings suggest that anxiety following

bereavement substantially contributes to one's level of social support during this pivotal time. Paired with findings that some older adults attempt to avoid anything that reminds them of their spouse in order to cope with spousal loss (Somhlaba and Wait, 2009), future research should examine social avoidance, withdrawal, and interpersonal rejections as potential mechanisms between anxiety and low emotional social support following spousal loss in older adults.

Building on the broad literature supporting the importance of social support following bereavement and in older adulthood (Stroebe et al., 2005; Dean et al., 1990), the current research also found that low emotional social support predicted symptoms of high depression two and a half years later. It is possible that the relationship between social support and later depression in bereaving older adults occurs due to those with low social support having insufficient social outlets to cope and process the loss of their spouse, and lead to feelings of loneliness, isolation (van Baarsen, 2002; Golden et al., 2009; Van Baarsen et al., 1999), and emotional isolation (van Baarsen, 2002). Thus, future research should examine loneliness, and emotional isolation as potential pathways between low social support and later depressive symptoms in widows.

The present findings also relate to coping with stressful life events. According to theory, critical life events require major readjustment. Intensity of stress depends on whether the demands of a situation exceed individuals' emotional coping resources (Lazarus and Folkman, 1984). Additionally, according to Holmes and Rahe (Holmes and Rahe, 1967), the death of a spouse is the most stressful life change event because it requires the most adaptive coping behaviors and is accompanied by the most psychological distress. One study found that 14-months after spousal bereavement, about a third of bereaved individuals had difficulties developing new intimate relationships, and the problem was particularly salient among older adults (Horowitz et al., 1997). A similar study found that 25 months later, 37% of widowers and 58% of widows reported difficulties developing new relationships (Schneider et al., 1996). Furthermore, number of family and friends in the social network and visits from family members tend to decline over the first year and half of bereavement (Utz et al., 2014). Our findings add to prior findings suggesting that higher anxiety may be related to such social support decline (Torgrud et al., 2004).

Our results support prior findings that elevated anxiety (Maccallum et al., 2015; Meier et al., 2013; Prigerson et al., 1996; Byrne and Raphael, 1997; Kreicbergs et al., 2004; Valdimarsdottir et al., 2004; Mitchell et al., 2009) and depressive symptoms (Kreicbergs et al., 2004; Mitchell et al., 2009; Stroebe et al., 1996, 2005; Murphy, 1988; Jozwiak et al., 2013; Ghesquiere et al., 2013) may hold particular importance following a loss, especially in regard to the social adjustment of the widow or widower. Additionally, social support may act as a protective factor against potentially adverse effects of stressful life events, as has been found previously; specifically, high levels of social support were associated with low depressive symptoms (Stroebe et al., 2005, 1996; Aneshensel and Frerichs, 1982; Cohen and Hoberman, 1983). Therefore, social support is particularly important for protecting against psychological distress following what is considered the most stressful life event, spousal bereavement. Prior studies also found that bereavement increased the need for social support (Stroebe and Stroebe, 1987). Additionally, those who had lost a spouse but perceived that they were more socially supported were less distressed than those who felt less socially supported (Bock and Webber, 1972; Dimond et al., 1987; Morgan, 1976). These results may suggest that interventions for bereaved older adults should include a component targeted toward reducing anxiety symptoms and fostering approach strategies, rather than avoidance strategies. They also suggest that targeting social support may be helpful to bereaved individuals, especially those exhibiting anxiety symptoms. In addition, because we found this relationship within the first 4 years' post-bereavement, it is possible that anxiety may rapidly progress toward later depressive symptoms.

³ Note that the model including the 11-item CES-D had good fit (PPP = 0.058). The pattern of emotional social support mediating the relationship between anxiety and depression held, as anxiety significantly negatively predicted emotional social support ($B = -0.291$, $CI = -0.586 - -0.012$, $d = -0.282$) and emotional social support significantly negatively predicted depression ($B = -0.306$, $CI = -0.502 - -0.123$, $d = -0.446$). Further the indirect effect of anxiety on depression, mediated by emotional social support, was also significantly positive ($B = 0.083$, $CI = 0.003 - 0.208$, $d = 0.221$). Additionally, the direct effect of anxiety on later depression was no longer significant ($B = 0.282$, $CI = -0.032 - 0.622$, $d = 0.237$). Also like the 6-item CES-D results, the results suggested that baseline depression did not significantly predict social support ($B = -0.119$, $CI = -0.390 - 0.149$, $d = -0.123$), and social support did not significantly predict later anxiety ($B = -0.017$, $CI = -0.220 - 0.175$, $d = -0.024$). The total effect of depression on anxiety ($B = 0.554$, $CI = 0.256 - 0.895$, $d = 0.480$) remained significant, but the indirect effect of depression on later anxiety mediated by social support ($B = 0.000$, $CI = -0.038 - 0.041$, $d = 0.000$) was not significant for the 11-item CES-D. Consequently, the theoretical implications of the model, including the reduced CES-D compared to the 11-item CES-D, are the same.

⁴ Note that the model including the stress covariates also had good fit (PPP = 0.058). Likewise, the pattern of emotional social support mediating the relationship between anxiety and depression held, as anxiety significantly negatively predicted emotional social support ($B = -0.333$, $CI = -0.665 - -0.050$, $d = -0.316$) and emotional social support significantly negatively predicted depression ($B = -0.633$, $CI = -0.872 - -0.442$, $d = -0.799$). Further the indirect effect of anxiety on depression, mediated by emotional social support, was also significantly positive ($B = 0.211$, $CI = 0.024 - 0.426$, $d = 0.301$). Additionally, the direct effect of anxiety on later depression was no longer significant ($B = 0.184$, $CI = -0.192 - 0.554$, $d = 0.140$). Likewise when controlling for stress covariates, the results suggested that baseline depression did not significantly predict social support ($B = -0.129$, $CI = -0.450 - 0.176$, $d = -0.121$), and social support did not significantly predict later anxiety ($B = -0.184$, $CI = -0.390 - 0.024$, $d = -0.248$). The total effect of depression on anxiety ($B = 0.367$, $CI = 0.080 - 0.633$, $d = 0.358$) remained significant, but the indirect effect of depression on later anxiety mediated by social support ($B = 0.016$, $CI = -0.031 - 0.101$, $d = 0.067$) was not significant when controlling for stress covariates. Consequently, the theoretical implications of the model, including the stress covariates, are the same.

Thus, intervention programs may be especially important during the first four years after loss of a spouse for older adults.

These results are consistent with and incrementally extend prior findings, as well as corroborate literature showing that anxiety and depressive symptoms bi-directionally predict one another years later (Jacobson and Newman, 2012b; van Baarsen, 2002). This is the first time a temporal mechanism between anxiety and depression has been examined in an older adult sample; all prior studies used adolescent or young adult samples. Nevertheless, in younger populations outside the context of bereavement, prior research found four constructs that partially longitudinally mediated the relationship between anxiety and depressive symptoms: avoidance, sociability, interpersonal oversensitivity, and perceptions of close and group relationships (Jacobson and Newman, 2014, 2016; Starr et al., 2014). The present findings add poor interpersonal relationships to this list, suggesting that perceived emotional support may be a mechanism by which anxiety predicts later depression. Furthermore, in pairing prior literature with the current results, the depth of one's social connections may be as influential as breadth in mediating the relationship between anxiety and later depression.

Although this study makes important contributions to the bereavement literature as well as the broader anxiety and depression literature, it is not without limitations. Additionally, the current research only utilized self-report measures, and, consequently, the results could be impacted by potential response biases. Consequently, the results should be replicated using multi-trait multimethod research with multiple informants. In particular, although we found a bi-directional longitudinal relationship between anxiety and depression following bereavement, we only found that social support unidirectionally mediated anxiety and later depression. Although little is known about the relationship between depression and later anxiety, future research should examine behavioral activation and avoidance as potential mediational candidates. In particular, depression was associated with low behavioral activation (Kasch et al., 2002). Decreases in behavioral activation could thereby decrease exposure to novel situations and experiences (Kasch et al., 2002) and further exacerbate fear and avoidance cycles (Kasch et al., 2002).

The current study sample also was not diverse, consisting primarily of heterosexual Caucasian females. Although a prior meta-analysis found that the relationship between anxiety and depressive symptoms was robust across gender and ethnicity (Jacobson and Newman, 2012b), more research is needed to determine whether the mediating role of social support applies across different gender identities, sexual orientations, and ethnicities. We were also unable to control for other third-variables that could have connections with the constructs examined here (i.e. avoidance, neuroticism, coping techniques, religiosity), and, consequently, future work should examine perceived emotional social support while also examining other potentially related control variables. Further, the current study used an emotional social support measure that has not been used in prior work, and, consequently, future work should attempt to replicate our findings with a validated measure. Moreover, although the study established emotional social support as mediating a large portion of the variance between anxiety and later depressive symptoms following bereavement, the current study could not establish mechanisms between anxiety and social support or between social support and depression. Likewise, although mediation provides a framework to examine mechanisms between processes, it should be noted that this does not imply that the statistical findings represent causal relationships, but rather a pattern of longitudinal findings that are indicative of risk factors. Notably, although the current use of structural equation modeling allows one to account for item-level measurement error from the examination of the relationship between constructs, it sacrifices the uniqueness of the importance of some items in favor of the shared covariance. Thus, future research should examine potential mediators between anxiety symptoms and social support, such as avoidance, and social support and depressive symptoms, such as loneliness.

Acknowledgements

The availability of these data is made possible by a grant from the National Institute on Aging (Randolph M. Nesse, Principal Investigator, AG15948-01). The original data collection for the CLOC study was supported by NIA grants (Camille B. Wortman, Principal Investigator, AG610757-01, and James S. House, Principal Investigator, AG05561-01).

References

- Aneshensel, C.S., Frerichs, R.R., 1982. Stress, support, and depression: A longitudinal causal model. *J. Community Psychol.* 10, 363–376. [http://dx.doi.org/10.1002/1520-6629\(198210\)10:4 < 363.](http://dx.doi.org/10.1002/1520-6629(198210)10:4 < 363.)
- Asparouhov, T., Muthen, B., 2010. Bayesian Analysis of Latent Variable Models using Mplus.
- Balaszamy, S., Richardson, V., Price, C.A., 2004. Investigating patterns of social support use by widowers during bereavement. *J. Men'S. Stud.* 13, 67–84. <http://dx.doi.org/10.3149/jms.1301.67.>
- Bennett, K.M., Smith, P.T., Hughes, G.M., 2005. Coping, depressive feelings and gender differences in late life widowhood. *Aging Ment. Health* 9, 348–353. <http://dx.doi.org/10.1080/13607860500089609.>
- Bergman, E.J., Haley, W.E., Small, B.J., 2010. The role of grief, anxiety, and depressive symptoms in the use of bereavement services. *Death Stud.* 34, 441–458. <http://dx.doi.org/10.1080/07481181003697746.>
- Bock, E.W., Webber, L.L., 1972. Suicide among the Elderly: Isolating Widowhood and Mitigating Alternatives. *J. Marriage Fam.* 34, 24–31. <http://dx.doi.org/10.2307/349626.>
- Boelen, P.A., Prigerson, H.G., 2007. The influence of symptoms of prolonged grief disorder, depression, and anxiety on quality of life among bereaved adults. *Eur. Arch. Psychiatry Clin. Neurosci* 257, 444–452. <http://dx.doi.org/10.1007/s00406-007-0744-0.>
- Bonanno, G.A., Mihalecz, M.C., LeJeune, J.T., 1999. The core emotion themes of conjugal loss. *Motiv. Emot.* 23, 175–201. <http://dx.doi.org/10.1023/a:1021398730909.>
- Broadhead, W.E., Gehlbach, S.H., de Gruy, F.V., Kaplan, B.H., 1988. The Duke-UNC Functional Social Support Questionnaire. Measurement of social support in family medicine patients. *Med. Care* 26, 709–723. <http://dx.doi.org/10.1097/00005650-198807000-00006.>
- Bush, B.A., Novack, T.A., Schneider, J.J., Madan, A., 2004. Depression following traumatic brain injury: The validity of the CES-D as a brief screening device. *J. Clin. Psychol. Med. Settings* 11, 195–201. <http://dx.doi.org/10.1023/B:JOC.0000037613.69367.d4.>
- Byrne, G.J., Raphael, B., 1997. The psychological symptoms of conjugal bereavement in elderly men over the first 13 months. *Int. J. Geriatr. Psychiatry* 12, 241–251. [http://dx.doi.org/10.1002/\(SICI\)1099-1166\(199702\)12:2 < 241::AID-GPS590 > 3.0.CO;2-0.](http://dx.doi.org/10.1002/(SICI)1099-1166(199702)12:2 < 241::AID-GPS590 > 3.0.CO;2-0.)
- Cacciatore, J., Schnebly, S., Froen, J.F., 2009. The effects of social support on maternal anxiety and depression after stillbirth. *Health Soc. Care Community* 17, 167–176. <http://dx.doi.org/10.1111/j.1365-2524.2008.00814.x.>
- Carnelley, K.B., Wortman, C.B., Bolger, N., Burke, C.T., 2006. The time course of grief reactions to spousal loss: evidence from a national probability sample. *J. Pers. Soc. Psychol.* 91, 476–492. <http://dx.doi.org/10.1037/0022-3514.91.3.476.>
- Chen, J.H., Bierhals, A.J., Prigerson, H.G., Kasl, S.V., Mazure, C.M., Jacobs, S., 1999. Gender differences in the effects of bereavement-related psychological distress in health outcomes. *Psychol. Med* 29, 367–380. <http://dx.doi.org/10.1017/s003291798008137.>
- Cohen, J., 1988. *Statistical power analysis for the behavioural sciences*. Lawrence Erlbaum Associates, NJ.
- Cohen, S., Hoberman, H.M., 1983. Positive events and social supports as buffers of life change stress. *J. Appl. Soc. Psychol.* 13, 99–125. <http://dx.doi.org/10.1111/j.1559-1816.1983.tb02325.x.>
- Colby, S.L., Ortman, J.M., 2015. Projections of the size and composition of the U.S. population: 2014 to 2060, Current Population Reports. Census Bureau, Washington DC: U.S..
- Condon, J.T., 1986. Management of established pathological grief reaction after stillbirth. *Am. J. Psychiatry* 143, 987–992. <http://dx.doi.org/10.1176/ajp.143.8.987.>
- Cornwell, E.Y., Waite, L.J., 2009. Social disconnectedness, perceived isolation, and health among older adults. *J. Health Soc. Behav.* 50, 31–48. <http://dx.doi.org/10.1177/002214650905000103.>
- Cumming, G., 2014. The new statistics: Why and how. *Psychol. Sci* 25, 7–29. <http://dx.doi.org/10.1177/0956797613504966.>
- Dean, A., Kolody, B., Wood, P., 1990. Effects of Social Support from Various Sources on Depression in Elderly Persons. *J. Health Soc. Behav.* 31, 148–161. <http://dx.doi.org/10.2307/2137169.>
- Derogatis, L.R., 1983. *SCL-90: Administration, scoring and procedures manual-I for the revised version and other instruments of the psychopathology rating scale series*. Johns Hopkins University School of Medicine, Clinical Psychometrics Research Unit, Baltimore, MD.
- Derogatis, L.R., Cleary, P.A., 1977a. Confirmation of the dimensional structure of the SCL-90: A study in construct validation. *J. Clin. Psychol.* 33, 981–989. <http://>

- [dx.doi.org/10.1002/1097-4679\(197710\)33:4 <81.1.110](http://dx.doi.org/10.1002/1097-4679(197710)33:4 <81.1.110)
- Derogatis, L.R., Cleary, P.A., 1977b. Factorial invariance across gender for the primary symptom dimensions of the SCL-90. *Br. J. Soc. Clin. Psychol* 16, 347–356. <http://dx.doi.org/10.1111/j.2044-8260.1977.tb00241.x>.
- Derogatis, L.R., Rickels, K., Rock, A.F., 1976. The SCL-90 and the MMPI: A step in the validation of a new self-report scale. *Br. J. Psychiatry* 128 (280), 280–289. <http://dx.doi.org/10.1192/bjp.128.3.280>.
- Dimond, M., Lund, D.A., Caserta, M.S., 1987. The role of social support in the first two years of bereavement in an elderly sample. *Gerontologist* 27, 599–604. <http://dx.doi.org/10.1093/geront/27.5.599>.
- Dunlap, W.P., Cortina, J.M., Vaslow, J.B., Burke, M.J., 1996. Meta-analysis of experiments with matched groups or repeated measures designs. *Psychol. Methods* 1, 170–177. <http://dx.doi.org/10.1037/1082-989X.1.2.170>.
- Dykstra, P.A., van Tilburg, T.G., Gierveld, J.J., 2005. Changes in older adult loneliness: Results from a seven-year longitudinal study. *Res. Aging* 27, 725–747. <http://dx.doi.org/10.1177/0164027505279712>.
- Enders, C.K., 2010. *Applied missing data analysis*. Guilford Press, New York, NY.
- Erickson, T.M., Newman, M.G., Siebert, E.C., Carlile, J.A., Scarsella, G.M., Abelson, J.L., 2016. Does worrying mean caring too much? Interpersonal prototypicality of dimensional worry controlling for social anxiety and depressive symptoms. *Behav. Ther.* 47, 14–28. <http://dx.doi.org/10.1016/j.beth.2015.08.003>.
- Field, N.P., Gal-Oz, E., Bonanno, G.A., 2003. Continuing bonds and adjustment at 5 years after the death of a spouse. *J. Consult. Clin. Psychol.* 71, 110–117. <http://dx.doi.org/10.1037/0022-006X.71.1.110>.
- Friedmann, E., Son, H., Thomas, S.A., Chapa, D.W., Lee, H.J., 2014. Poor social support is associated with increases in depression but not anxiety over 2 years in heart failure outpatients. *J. Cardiovasc. Nurs.* 29, 20–28. <http://dx.doi.org/10.1097/JCN.0b013e318276fa07>.
- Galatzer-Levy, I.R., Bonanno, G.A., 2012. Beyond normality in the study of bereavement: heterogeneity in depression outcomes following loss in older adults. *Soc. Sci. Med.* 74, 1987–1994. <http://dx.doi.org/10.1016/j.socscimed.2012.02.022>.
- Gellis, Z.D., 2010. Assessment of a brief CES-D measure for depression in homebound medically ill older adults. *J. Gerontol. Soc. Work* 53, 289–303. <http://dx.doi.org/10.1080/01634371003741417>.
- Gelman, A., Rubin, D.B., 1992. Inference from iterative simulation using multiple sequences. *Stat. Sci.* 7, 457–472. <http://dx.doi.org/10.1214/ss/1177011136>.
- Ghesquiere, A., Shear, M.K., Duan, N., 2013. Outcomes of bereavement care among widowed older adults with complicated grief and depression. *J. Prim. Care Community Health* 4, 256–264. <http://dx.doi.org/10.1177/2150131913481231>.
- Golden, J., Conroy, R.M., Bruce, I., Denihan, A., Greene, E., Kirby, M., Lawlor, B.A., 2009. Loneliness, social support networks, mood and wellbeing in community-dwelling elderly. *Int. J. Geriatr. Psychiatry* 24, 694–700. <http://dx.doi.org/10.1002/gps.2181>.
- Hansson, R.O., Stroebe, M.S., 2007. *Bereavement in late life: Coping, adaptation, and developmental influence*. s. American Psychological Association, Washington, DC, US.
- Henderson, S., Duncan-Jones, P., Byrne, D.G., Scott, R., 1980. Measuring social relationships. The Interview Schedule for Social Interaction. *Psychol. Med* 10, 723–734. <http://dx.doi.org/10.1017/S003329170005501X>.
- Hogan, N., Morse, J.M., Tasón, M.C., 1996. Tasón Toward an experiential theory of bereavement. *J. Death and Dying* 33, 43–65. <http://dx.doi.org/10.2190/gu3x-jwv0-ag6g-21fx>.
- Holland, J.M., Thompson, K.L., Rozalski, V., Lichtenthal, W.G., 2014. Bereavement-related regret trajectories among widowed older adults. *J. Gerontol. B. Psychol. Sci. Soc.* 69, 40–47. <http://dx.doi.org/10.1093/geronb/gbt050>.
- Holmes, T.H., Rahe, R.H., 1967. The social readjustment rating scale. *J. Psychosom. Res.* 11, 213–218. [http://dx.doi.org/10.1016/0022-3999\(67\)90010-4](http://dx.doi.org/10.1016/0022-3999(67)90010-4).
- Horowitz, M.J., Siegel, B., Holen, A., Bonanno, G.A., Milbrath, C., Stinson, C.H., 1997. Diagnostic criteria for complicated grief disorder. *Am. J. Psychiatry* 154, 904–910. <http://dx.doi.org/10.1176/ajp.154.7.904>.
- Jacobson, N.C., Newman, M.G., 2012a. Anxiety as a temporal predictor of depression during brief time periods: An ecological momentary assessment November 2012, National Harbor. Association for Behavioral and Cognitive Therapies, National Harbor, MD, Maryland, 18.
- Jacobson, N.C., Newman, M.G., 2012b. The temporal relationship between anxiety and depression: A meta-analysis November 2012, National Harbor. Association for Behavioral and Cognitive Therapies National Harbor, MD, Maryland, 18.
- Jacobson, N.C., Newman, M.G., 2014. Avoidance mediates the relationship between anxiety and depression over a decade later. *J. Anxiety Disord* 28, 437–445. <http://dx.doi.org/10.1016/j.janxdis.2014.03.007>.
- Jacobson, N.C., Newman, M.G., 2016. Perceptions of close and group relationships mediate the relationship between anxiety and depression over a decade later. *Depress. Anxiety* 33, 66–74. <http://dx.doi.org/10.1002/da.22402>.
- Jacobson, N.C., Newman, M.G., 2017. M.G. Anxiety and depression as bidirectional risk factors for one another: A meta-analysis of longitudinal studies. (in preparation)
- Jaggal, H.S., 1982. Multicollinearity in structural equation models with unobservable variables. *J. Mark. Res.* 19, 431.
- Jensen, M.P., Smith, A.E., Bombardier, C.H., Yorkston, K.M., Miro, J., Molton, I.R., 2014. Social support, depression, and physical disability: Age and diagnostic group effects. *Disabil. Health J* 7, 164–172. <http://dx.doi.org/10.1016/j.dhjo.2013.11.001>.
- Jozwiak, N., Preville, M., Vasiladias, H.M., 2013. Bereavement-related depression in the older adult population: A distinct disorder? *J. Affect. Disord* 151, 1083–1089. <http://dx.doi.org/10.1016/j.jad.2013.08.038>.
- Kahn, J.H., Garrison, A.M., 2009. Emotional self-disclosure and emotional avoidance: Relations with symptoms of depression and anxiety. *J. Couns. Psychol.* 56, 573–584. <http://dx.doi.org/10.1037/a0016574>.
- Kasch, K.L., Rottenberg, J., Arnow, B.A., Gotlib, I.H., 2002. Behavioral activation and inhibition systems and the severity and course of depression. *J. Abnorm. Psychol.* 111, 589–597. <http://dx.doi.org/10.1037/0021-843X.111.4.589>.
- Kass, R.E., Raftery, A.E., 1995. Bayes Factors. *J. Am. Stat. Assoc.* 90, 773–795. <http://dx.doi.org/10.1080/01621459.1995.10476572>.
- Keltner, D., Bonanno, G.A., 1997. A study of laughter and dissociation: distinct correlates of laughter and smiling during bereavement. *J. Pers. Soc. Psychol.* 73, 687–702. <http://dx.doi.org/10.1037/0022-3514.73.4.687>.
- Kohout, F., 1992. The pragmatics of survey field work among the elderly. In: Wallace, R., Woolson, R. (Eds.), *The epidemiological study of the elderly*. Oxford University Press, New York, 99–119.
- Kreicbergs, U., Valdimarsdottir, U., Onelov, E., Henter, J.I., Steineck, G., 2004. Anxiety and depression in parents 4–9 years after the loss of a child owing to a malignancy: A population-based follow-up. *Psychol. Med* 34, 1431–1441. <http://dx.doi.org/10.1017/s0033291704002740>.
- Langston, C.A., Cantor, N., 1989. Social anxiety and social constraint: When making friends is hard. *J. Pers. Soc. Psychol.* 56, 649–661. <http://dx.doi.org/10.1037/0022-3514.56.4.649>.
- Lazarus, R.S., Folkman, S., 1984. *Stress, appraisal and the coping process*. Springer, New York.
- Maccallum, F., Sawday, S., Rinck, M., Bryant, R.A., 2015. The push and pull of grief: Approach and avoidance in bereavement. *J. Behav. Ther. Exp. Psychiatry* 48, 105–109. <http://dx.doi.org/10.1016/j.jbtep.2015.02.010>.
- Meier, A.M., Carr, D.R., Currier, J.M., Neimeyer, R.A., 2013. Attachment anxiety and avoidance in coping with bereavement: Two studies. *J. Soc. Clin. Psychol.* 32, 315–334. <http://dx.doi.org/10.1521/jscp.2013.32.3.315>.
- Middleton, W., Burnett, P., Raphael, B., Martinek, N., 1996. The bereavement response: A cluster analysis. *Br. J. Psychiatry* 169, 167–171. <http://dx.doi.org/10.1192/bjp.169.2.167>.
- Mitchell, A.M., Sakranda, T.J., Kim, Y., Bullian, L., Chiappetta, L., 2009. Depression, anxiety and quality of life in suicide survivors: A comparison of close and distant relationships. *Arch. Psychiatr. Nurs* 23, 2–10. <http://dx.doi.org/10.1016/j.apnu.2008.02.007>.
- Morgan, L.A., 1976. A re-examination of widowhood and morale. *J. Gerontol* 31, 687–695.
- Mullins, L.C., Dugan, E., 1990. The influence of depression, and family and friendship relations, on residents' loneliness in congregate housing. *Gerontologist* 30, 377–384.
- Murphy, S.A., 1988. Mental distress and recovery in a high-risk bereavement sample three years after untimely death. *Nurs. Res* 37, 30–35. <http://dx.doi.org/10.1097/00006199-198801000-00007>.
- Muthén, B., Asparouhov, T., 2012a. Bayesian structural equation modeling: A more flexible representation of substantive theory. *Psychol. Methods* 17, 313–335. <http://dx.doi.org/10.1037/a0026802>.
- Muthén, B., Asparouhov, T., 2012b. Rejoinder to MacCallum, Edwards, and Cai (2012) and Rindskopf (2012): Mastering a new method. *Psychol. Methods* 17, 346–353. <http://dx.doi.org/10.1037/a0029214>.
- Nesse, R.M., Wortman, C., House, J., Kessler, R., Lepkowski, J., 2006. Changing lives of older couples (CLOC): A study of spousal bereavement in the Detroit area, 1987–1993. *Inter-Univ. Consort. Political Soc. Res.* <http://dx.doi.org/10.3886/ICPSR03370.v1>.
- Newcombe, R.G., 1998. Two-sided confidence intervals for the single proportion: Comparison of seven methods. *Stat. Med* 17, 857–872.
- Ong, A.D., Bergeman, C.S., Bisconti, T.L., 2005. Unique effects of daily perceived control on anxiety symptomatology during conjugal bereavement. *Personal. Individ. Differ.* 38, 1057–1067. <http://dx.doi.org/10.1016/j.paid.2004.07.004>.
- Paterson, C., Robertson, A., Nabi, G., 2015. Exploring prostate cancer survivors' self-management behaviours and examining the mechanism effect that links coping and social support to health-related quality of life, anxiety and depression: A prospective longitudinal study. *Eur. J. Oncol. Nurs.* 19, 120–128. <http://dx.doi.org/10.1016/j.ejon.2014.10.008>.
- Post, A.L., Wittmaier, B.C., Radin, M.E., 1978. Self-disclosure as a function of state and trait anxiety. *J. Consult. Clin. Psychol.* 46, 12–19. <http://dx.doi.org/10.1037/0022-006X.46.1.12>.
- Prigerson, H.G., Shear, M.K., Newsom, J.T., Frank, E., Reynolds, C.F., Maciejewski, P.K., Houck, P.R., Bierhals, A.J., Kupfer, D.J., 1996. Anxiety among widowed elders: Is it distinct from depression and grief? *Anxiety* 2, 1–12. [http://dx.doi.org/10.1002/\(sici\)1522-7154\(1996\)2:1 <1::aid-anx1 >3.3.co;2-2](http://dx.doi.org/10.1002/(sici)1522-7154(1996)2:1 <1::aid-anx1 >3.3.co;2-2).
- Prince, M.J., Harwood, R.H., Blizard, R.A., Thomas, A., Mann, A.H., 1997. Social support deficits, loneliness and life events as risk factors for depression in old age. *The Gospel Oak Project VI. Psychol. Med* 27, 323–332.
- Radloff, L.S., 1977. The CES-D Scale: A self-report depression scale for research in the general population. *Appl. Psych. Meas.* 1, 385–401. <http://dx.doi.org/10.1177/014662167700100306>.
- Rupp, A.A., Dey, D.K., Zumbo, B.D., 2004. To Bayes or not to Bayes, from whether to when: Applications of Bayesian methodology to modeling. *Structural Equ. Model.: A Multidiscip. J.* 11, 424–451. http://dx.doi.org/10.1207/s15328007sem1103_7.
- Sasson, I., Umberson, D.J., 2014. Widowhood and depression: New light on gender differences, selection, and psychological adjustment. *J. Gerontol. Ser. B: Psychol. Sci. Soc. Sci.* 69B, 135–145. <http://dx.doi.org/10.1093/geronb/gbt058>.
- Scharfstein, L., Alfano, C., Beidel, D., Wong, N., 2011. Children with generalized anxiety disorder do not have peer problems, just fewer friends. *Child Psychiatry Hum. Dev.* 42, 712–723. <http://dx.doi.org/10.1007/s10578-011-0245-2>.
- Schneider, D.S., Sledge, P.A., Shuchter, S.R., Zisook, S., 1996. Dating and remarriage over the first two years of widowhood. *Ann. Clin. Psychiatry* 8, 51–57. <http://dx.doi.org/10.3109/10401239609148802>.
- Sherbourne, C.D., Stewart, A.L., 1991. The MOS social support survey. *Soc. Sci. Med.* 32,

- 705–714. [http://dx.doi.org/10.1016/0277-9536\(91\)90150-B](http://dx.doi.org/10.1016/0277-9536(91)90150-B).
- Sibbritt, D.W., Byles, J.E., Regan, C., 2007. Factors associated with decline in physical functional health in a cohort of older women. *Age Ageing* 36, 382–388. <http://dx.doi.org/10.1093/ageing/afm017>.
- Somhlaba, N.Z., Wait, J.W., 2009. Stress, coping styles, and spousal bereavement: Exploring patterns of grieving among black widowed spouses in rural South Africa. *J. Loss Trauma: Int. Perspect. Stress Coping* 14, 192–210. <http://dx.doi.org/10.1080/15325020802537443>.
- Starr, L.R., Davila, J., 2012a. Cognitive and interpersonal moderators of daily co-occurrence of anxious and depressed moods in generalized anxiety disorder. *Cognit. Ther. Res* 36, 655–669. <http://dx.doi.org/10.1007/s10608-011-9434-3>.
- Starr, L.R., Davila, J., 2012b. Temporal patterns of anxious and depressed mood in generalized anxiety disorder: A daily diary study. *Behav. Res. Ther* 50, 131–141. <http://dx.doi.org/10.1016/j.brat.2011.11.005>.
- Starr, L.R., Hammen, C., Connolly, N.P., Brennan, P.A., 2014. Does relational dysfunction mediate the association between anxiety disorders and later depression? Testing an interpersonal model of comorbidity. *Depress. Anxiety* 31, 77–86. <http://dx.doi.org/10.1002/da.22172>.
- Starr, L.R., Stroud, C.B., Li, Y.I., 2016. Predicting the transition from anxiety to depressive symptoms in early adolescence: Negative anxiety response style as a moderator of sequential comorbidity. *J. Affect. Disord* 190, 757–763. <http://dx.doi.org/10.1016/j.jad.2015.10.065>.
- Steer, R.A., Ranieri, W.F., Beck, A.T., Clark, D.A., 1993. Further evidence for the validity of the Beck Anxiety Inventory with psychiatric outpatients. *J. Anxiety Disord* 7, 195–205. [http://dx.doi.org/10.1016/0887-6185\(93\)90002-3](http://dx.doi.org/10.1016/0887-6185(93)90002-3).
- Stekhoven, D.J., Bühlmann, P., 2012. MissForest—non-parametric missing value imputation for mixed-type data. *Bioinformatics* 28, 112–118. <http://dx.doi.org/10.1093/bioinformatics/btr597>.
- Stroebe, W., Stroebe, M., Abakoumkin, G., Schut, H., 1996. The role of loneliness and social support in adjustment to loss: A test of attachment versus stress theory. *J. Pers. Soc. Psychol.* 70, 1241–1249. <http://dx.doi.org/10.1037//0022-3514.70.6.1241>.
- Stroebe, W., Stroebe, M.S., 1987. *Bereavement and health: The psychological and physical consequences of partner loss*. Cambridge University Press, New York, NY.
- Stroebe, W., Zech, E., Stroebe, M.S., Abakoumkin, G., 2005. Does social support help in bereavement? *J. Soc. Clin. Psychol.* 24, 1030–1050. <http://dx.doi.org/10.1521/jscp.2005.24.7.1030>.
- Swendsen, J.D., 1997. Anxiety, depression, and their comorbidity: An experience sampling test of the helplessness-hopelessness theory. *Cognit. Ther. Res* 21, 97–114. <http://dx.doi.org/10.1023/A:1021872410824>.
- Team, J. (Ed.), 2016. *JASP* 0.8.
- Torgrud, L.J., Walker, J.R., Murray, L., Cox, B.J., Chartier, M., Kjernisted, K.D., 2004. Deficits in perceived social support associated with generalized social phobia. *Cogn. Behav. Ther* 33, 87–96. <http://dx.doi.org/10.1080/16506070410029577>.
- Uden, A.L., Orth-Gomer, K., 1989. Development of a social support instrument for use in population surveys. *Soc. Sci. Med* 29, 1387–1392. [http://dx.doi.org/10.1016/0277-9536\(89\)90240-2](http://dx.doi.org/10.1016/0277-9536(89)90240-2).
- United States Census Bureau, 2015. *America's Families and Living Arrangements, 2015, Adults*.
- Utz, R.L., Caserta, M., Lund, D., 2011. Grief, depressive symptoms, and physical health among recently bereaved spouses. *Gerontologist* 52, 460–471. <http://dx.doi.org/10.1093/geront/gnr110>.
- Utz, R.L., Swenson, K.L., Caserta, M., Lund, D., deVries, B., 2014. Feeling lonely versus being alone: loneliness and social support among recently bereaved persons. *J. Gerontol B Psychol Sci Soc Sci* 69B, 85–94. <http://dx.doi.org/10.1093/geronb/gbt075>.
- Valdimarsdottir, U., Helgason, A.R., Furst, C.J., Adolffson, J., Steineck, G., 2004. Awareness of husband's impending death from cancer and long-term anxiety in widowhood: A nationwide follow-up. *Palliat. Med* 18, 432–443. <http://dx.doi.org/10.1191/0269216304pm891oa>.
- van Baarsen, B., 2002. Theories on coping with loss: The impact of social support and self-esteem on adjustment to emotional and social loneliness following a partner's death in later life. *J. Gerontol. B. Psychol. Sci. Soc. Sci* 57, S33–42.
- van Baarsen, B., Smit, J.H., Snijders, T.A.B., Knipscheer, K.P.M., 1999. Do personal conditions and circumstances surrounding partner loss explain loneliness in newly bereaved older adults? *Ageing Soc.* 19, 441–469.
- van Grootheest, D.S., Beekman, A.T., Broese van Groenou, M.I., Deeg, D.J., 1999. Sex differences in depression after widowhood. Do men suffer more? *Soc. Psychiatry Psychiatr. Epidemiol.* 34, 391–398. <http://dx.doi.org/10.1007/s001270050160>.
- Victor, C., Scambler, S., Bond, J., Bowling, A., 2000. Being alone in later life: Loneliness, social isolation and living alone. *Rev. Clin. Gerontol.* 10, 407–417.
- Ward, L., Mathias, J.L., Hitchings, S.E., 2007. Relationships between bereavement and cognitive functioning in older adults. *Gerontology* 53, 362–372. <http://dx.doi.org/10.1159/000104787>.
- Weissman, M.M., Sholomskas, D., Pottenger, M., Prusoff, B.A., Locke, B.Z., 1977. Assessing depressive symptoms in five psychiatric populations: A validation study. *Am. J. Epidemiol* 106, 203–214.
- Williams, B.R., Sawyer Baker, P., Allman, R.M., Roseman, J.M., 2007. Bereavement among African American and White older adults. *J. Aging Health* 19, 313–333. <http://dx.doi.org/10.1177/0898264307299301>.
- Winship, C., 1999. *The perils of multicollinearity: A reassessment*. Harvard University.
- Yuan, Y., MacKinnon, D.P., 2009. Bayesian mediation analysis. *Psychol. Methods* 14, 301–322. <http://dx.doi.org/10.1037/a0016972>.